

1/20




FIG. 1

09839577.082001

2/20




FIG. 2

3/20




FIG. 3

4/20




FIG. 4

5/20




FIG. 5

09839577 256860




FIG. 6A




FIG. 6B

7/20

GFP-Rac to Alexa-PBD FRET




FIG. 7A

8/20

FRET response to nucleotide state of Rac-GFP




FIG. 7B

9/20




FIG. 8A




FIG. 8B

11/20

FIG. 9A

SERUM STIMULATION OF A SWISS 3T3 FIBROBLAST




FIG. 9A

FIG. 9B

TITLE: LABELED PEPTIDES,

INS AND ANTIBODIES AND PROCESSES
FOR THEIR PREPARATION
INVENTORS NAME: Klaus M. Hahn et al.
SERIAL NO.: 09/839,577

INTERMEDIATES USEFUL

12/20

FIG. 9C

THE SAME RUFFLE VISUALIZED USING EITHER FRET OR Alexa-PBD LOCALIZATION:




FIG. 9D

FIG. 9C

FIG. 10A: Rac-GFP

Fig. 10A: Rac-GFP



Wound healing

Fig. 10B. FRET



Confluent
monolayer



Magnitude of gradient when highest at front	128 +/- 51 %	n=12
Magnitude of gradient highest at rear	9 +/- 4 %	n=4

14/20




FIG. 11A




FIG. 11B

15/20




FIG. 12A




FIG. 12B

Wavelength, nm

09839577 2256E860



17/20

FIG. 14




FIG. 14

18/20

Fluorescence of Mero-CBD responds to Cdc42 binding




FIG. 15

19/20




FIG. 16




FIG. 17